



Basis for Position on Mandatory Spay-Neuter in the Canine and Feline

The American College of Theriogenologists (ACT) is the certifying college for veterinarians board certified in reproduction (specialists) and the Society for Theriogenology (SFT) is an organization of veterinarians with a special interest in reproduction in veterinary medicine. **The ACT and SFT believe that companion animals not intended for breeding should be spayed or neutered; however, both organizations believe that the decision to spay or neuter a pet must be made on a case by case basis, taking into consideration the pet's age, breed, sex, intended use, household environment and temperament. The use of generalized rules concerning gonadectomy (removal of the ovaries or testes) is not in the best interest of the health or well-being of the pets or their owners.** Each of the following considerations must be assessed for each individual animal and household.

1) Health concerns

- a. Research has shown that there can be positive effects of the sex steroid hormones. The sex steroids are hormones produced by the ovaries and testes, and are only present in intact males and females. Gonadectomy at any age deprives the body of the positive health effects of these hormones. Although in most cases, the benefits of spay-neuter outweigh the benefits of exposure to the sex steroids, this is not true in all cases. Since gonadectomy prior to puberty or sexual maturity may make the risks of some diseases higher in certain breeds or individuals, the option to leave an animal intact must be available to the pet owner.
 - i. Advantages of remaining intact:
 1. There is a decreased incidence of hemangiosarcoma in intact bitches and dogs.
 2. There is a decreased incidence of osteosarcoma in intact male and female dogs.
 3. There is a decreased risk of transitional cell carcinoma in intact dogs and bitches.
 4. There is a decreased risk of prostatic adenocarcinoma in intact male dogs compared to gonadectomized male dogs.
 5. There is a decreased incidence of obesity in intact male and female dogs and cats, which may be due at least partly to increased metabolic rate.
 6. There is a decreased incidence of urinary incontinence in intact bitches (equivocal if bitches are spayed after 5 months but before their first heat).
 7. There may be a reduced incidence of urinary tract infection in intact bitches.
 8. There may be a reduced incidence of feline lower urinary tract disease (FLUTD) in intact male and female cats which may be partly due to decreased obesity in these animals.
 9. There may be a reduced incidence of autoimmune thyroiditis and hypothyroidism in intact male and female dogs.
 10. There is a decreased incidence of diabetes mellitus in intact female cats and a possibly reduced incidence in diabetes mellitus in intact male dogs.
 11. There is a reduced incidence of cranial cruciate rupture in intact male and female dogs.
 12. There may be a reduced incidence of hip dysplasia in male and female dogs that are not gonadectomized before 5 months of age.
 13. There may be an increased incidence of capital physeal fractures in castrated male cats that may be partially due to increased weight gain in gonadectomized males.
- b. Research has shown that there are a number of detrimental effects of the sex steroid hormones. Spaying and neutering will remove these hormones and thus lower the risk of these conditions.
 - i. Advantages of being spayed or castrated:

1. There is an increased risk of mammary, testicular, and ovarian neoplasia in intact male and female dogs and cats.
 - a. There is an increased risk of mammary cancer with each subsequent cycle and the benefit of spaying does not disappear until the animal reaches old age.
 - i. Mammary cancer is one of the most common types of neoplasia in small animals.
 1. Mammary neoplasia is malignant 60% of the time in dogs and 90% of the time in cats.
 - b. The incidence and mortality risk for ovarian cancer are very low
 - c. The incidence for testicular cancer is more common but malignancy and mortality are very low.
 2. There is an increased risk of pyometra in both intact female dogs and cats and this risk increases with increasing age.
 3. There is an increased risk of prostatitis, benign prostatic hyperplasia, prostatic cysts and squamous metaplasia of the prostate in intact male dogs.
 4. There is a decreased incidence of perineal and inguinal hernia and perineal adenoma in neutered male dogs.
- c. **Based on the research available, it is clear there are a number of health benefits of the sex steroid hormones and that this benefit varies with age, sex, and breed. Therefore, although spay-neuter is the responsible choice for most pets, it is in the best interest of each individual patient for its veterinarian to assess the risks and benefits of gonadectomy and to advise his/her clients on what is appropriate for each individual pet at each stage of its life.**
- 2) Behavioral concerns
- a. Research has shown that there are positive effects of the sex steroid hormones on behavior.
 - i. Advantages of remaining intact:
 1. There is a decrease in shyness and hiding behavior in intact male and female cats.
 2. There may be less aggression towards people and animals in intact bitches.
 3. There may be a decreased incidence of cognitive dysfunction in intact male and female dogs.
 - b. Research has shown that there are negative effects of the sex steroid hormones on behavior.
 - i. Advantages of being spayed or castrated:
 1. Inter-dog aggression may be due to competition for available territory or availability of cycling animals.
 2. Urine spraying and inter-animal aggression is increased in intact male cats.
 3. There is a decreased risk of wandering and being hit by a car in neutered animals.
- 3) Provision of quality medical care
- a. It is not in the animals' best interest to have the legislature dictate the time or need for surgical treatment.
 - i. This does not allow for medical decisions based on the individual animal's needs, its owners' needs or the needs of the household.
 1. Animals with medical conditions that may result in complications during anesthesia or surgery (i.e. heart murmurs, bleeding disorders).
 2. Providing appropriate aftercare for surgical patients may not be feasible in some home situations.
 - b. Restricting and reducing the pool of purebred animals will greatly hinder medical research of conditions that are particular to specific breeds, slowing down advances in medical and surgical knowledge. This may in turn impact the research available concerning health conditions common to both animals and people.
- 4) Public Health concerns
- a. Making spay/neuter mandatory for licensure may make the public more hesitant to seek veterinary assistance because they are afraid of fines and legal repercussions as a result of failing to spay or neuter their pets by the prescribed time. By avoiding veterinary care for their pets, animals will be at increased risk of inadequate routine vaccination (including rabies) and

inadequate deworming programs which may in turn result in increased transmission of disease to the public.

The ACT and SFT make the following recommendations to continue moving toward effective methods of reducing the number of abandoned, unwanted and euthanized dogs and cats in the US and other countries where similar problems exist.

- a. Provide increased jurisdictional control to the AVMA Governmental Relations division, Animal Welfare Committee, and the APHIS-Animal Care division.
- b. Ensure suppliers to pet stores are providing adequate care for breeding stock and offspring.
- c. Support programs to expand the public awareness of pet overpopulation, acceptable breeding standards, and responsibilities of pet ownership. Provide the public a means to access assistance with concerns of pet health, ownership, behavior and management issues.
- d. Work with state and local rescue and humane societies to assemble accurate data on causes for relinquishment of dogs and cats to enable these organizations, federal and local governments, and veterinary organizations to address the fundamental causes of abandonment.
- e. Provide low cost spay/neuter facilities for economically disadvantaged persons and communities.
- f. Continue to work on reduction of feral cat populations.
- g. Establish programs to ensure access of breeders to proper reproductive care and counseling.
- h. Provide local or federal governmental assistance to registered rescue organizations to facilitate placement of unwanted pets.

The ACT and SFT do not believe that **mandatory** spay/neuter programs will significantly reduce the pet overpopulation problems, since most animals that are abandoned are relinquished because of behavior, health, economic and life changing conditions and not due to their reproductive status. In fact, in some European Union countries where gonadectomy is illegal unless deemed medically necessary (such as Norway) there are no significant problems with pet overpopulation, indicating that the pet overpopulation problem that exists in the United States is due to cultural differences on the importance of pets, the responsibility of pet owners, and the ability of the government and national agencies to properly educate the public. Although both organizations believe that most companion animals should be spayed or neutered, the ACT and SFT also strongly believe that it is not in the best interest of the animals to produce legislation regarding medical treatments. Therefore, both organizations oppose **mandatory** spay/neuter programs.

There are hundreds of references which provide scientific information on the effects of spay and neuter in both dogs and cats. We chose to provide the reader with a selected list of them. This reference list was compiled by Dr. Peggy Root-Kustritz, DACT.

References:

- 1) Alexander SA, Shane SM. Characteristics of animals adopted from an animal control center whose owners complied with a spaying/neutering program. *J Amer Vet Med Assoc* 1994;205:472-276.
- 2) Angioletti A, DeFrancesco I, Vergottini M, et al. Urinary incontinence after spaying in the bitch: Incidence and oestrogen therapy. *Vet Res Commun* 2004;28 (Suppl 1): 153-155.
- 3) Arendt EA. Orthopaedic issues for active and athletic women. *Clin Sports Med* 1994;13:483-503.
- 4) Arnold S. Urinary incontinence in castrated bitches. Part I. Significance, clinical aspects and etiopathogenesis. *Schweiz Arch Tierheilkd* 1997;139:271-276.
- 5) Atalan G, Holt PE, Barr FJ. Ultrasonographic assessment of bladder neck mobility in continent bitches and bitches with urinary incontinence attributable to urethral sphincter mechanism incompetence. *Amer J Vet Res* 1998;59:673-679.
- 6) Beijerink NJ, Buijtels JJCWM, Okkens AC, et al. Basal and GnRH-induced secretion of FSH and LH in anestrus versus ovariectomized bitches. *Theriogenology* 2007;67:1039-1045.
- 7) Bell FW, Klausner JS, Hayden DW, et al. Clinical and pathologic features of prostatic adenocarcinoma in sexually intact and castrated dogs: 31 cases (1970-1987). *J Amer Vet Med Assoc* 1991;199:1623-1630.

- 8) Berry SJ, Strandberg JD, Saunders WJ, Coffey DS. Development of canine benign prostatic hyperplasia with age. *Prostate* 1986;9:363-373.
- 9) Brodey RS. Canine and feline neoplasia. *Adv Vet Sci Comp Med* 1970;14:309-354.
- 10) Brodey RS, Goldschmidt MH, Roszel JR. Canine mammary gland neoplasms. *J Amer Anim Hosp Assoc* 1983;19:61-90.
- 11) Bronson RT. Variation in age at death of dogs of different sexes and breeds. *Amer J Vet Res* 1982;43:2057-2059.
- 12) British Small Animal Veterinary Association. Sequelae of bitch sterilisation: Regional survey. *Vet Rec* 1975;96:371-372.
- 13) Bryan JN, Keeler MR, Henry CJ, et al. A population study of neutering status as a risk factor for canine prostate cancer. *Prostate* 2007;67:1174-1181.
- 14) Burrow R, Batchelor D, Cripps P. Complications observed during and after ovariohysterectomy of 142 bitches at a veterinary teaching hospital. *Vet Rec* 2005;157:829-833.
- 15) Coffey DJ. Sexual mutilation. *Vet Times*, Surrey UK, December 1998.
- 16) Cohen D, Reif JS, Brodey RS, Keiser H. Epidemiological analysis of the most prevalent sites and types of canine neoplasia observed in a veterinary hospital. *Cancer Res* 1974;34:2859-2868.
- 17) Cooley DM, Beranek BC, Schlittler DL, et al. Endogenous gonadal hormone exposure and bone sarcoma risk. *Canc Epidem Biomark Prev* 2002;11:1434-1440.
- 18) Cotchin E. Neoplasia in the dog. *Vet Rec* 1954;66:879-888.
- 19) Cotchin E. Neoplasms in small animals. *Vet Rec* 1951;63:67-72.
- 20) Cowan LA, Barsanti JA, Crowell W, Brown J. Effects of castration on chronic bacterial prostatitis in dogs. *J Amer Vet Med Assoc* 1991;199:346-350.
- 21) Crane SW. Occurrence and management of obesity in companion animals. *J Sm Anim Prac* 1991;32:275-282.
- 22) Crenshaw WE, Carter CN. Should dogs in animal shelters be neutered early? *Vet Med* 1995;90:756-760.
- 23) Daniels R. Personal communication, 03-20-2002.
- 24) DeSapio A, VanDeCoevering P, Williams JD, et al. Effects of early-age gonadectomy on a male guide dog population. Proceedings, International Working Dog Breed Conference, Sydney, Australia, 2005.
- 25) Donnay I, Raus J, Devleeshouwer N, et al. Comparison of estrogen and progesterone receptor expression in normal and tumor mammary tissues from dogs. *Amer J Vet Res* 1995;56:1188-1194.
- 26) Dorfman M, Barsanti J. Diseases of the canine prostate gland. *Comp Cont Ed* 1995;17:791-810.
- 27) Dorn CR, Taylor DON, Schneider R, et al. Survey of animal neoplasms in Alameda and Contra Costa counties, California. II. Cancer morbidity in dogs and cats from Alameda county. *J Nat Canc Inst* 1968;40:307-318.
- 28) Dorn CR, Taylor DON, Frye FL, Hibbard HH. Survey of animal neoplasms in Alameda and Contra Costa counties, California. I. Methodology and description of cases. *J Nat Canc Inst* 1968;40:295-305.

- 29) Duerr FM, Duncan CG, Savicky RS, et al. Risk factors for excessive tibial plateau angle in large-breed dogs with cranial cruciate ligament disease. *J Amer Vet Med Assoc* 2007;231:1688-1691.
- 30) Durham SK, Dietze AE. Prostatic adenocarcinoma with and without metastases to bone in dogs. *J Amer Vet Med Assoc* 1986;188:1432-1436.
- 31) Duval JM, Budsberg SC, Flo GL, Sammarco JL. Breed, sex, and body weight as risk factors for rupture of the cranial cruciate ligament in young dogs. *J Amer Vet Med Assoc* 1999;215:811-814.
- 32) Edney ATB, Smith PM. Study of obesity in dogs visiting veterinary practices in the United Kingdom. *Vet Rec* 1986;118:391-396.
- 33) Egenvall A, Hagman R, Bonnett BN, et al. Breed risk of pyometra in insured dogs in Sweden. *J Vet Intern Med* 2001;15:530-538.
- 34) Ekici H, Sontas BH, Toydemir TSF, et al. The effect of prepubertal ovariohysterectomy on spine 1 mineral density and mineral content in puppies: A preliminary study. *Res Vet Sci* 2007;82:105-109.
- 35) Eno M, Fekety S. Early age spay/neuter: A growing consensus. *Shelter Sense* 1993;Nov:1-7.
- 36) Eze CA, Eze MC. Castration, other management practices and socio-economic implications for dog keepers in Nsukka area, Enugu State, Nigeria. *Prev Vet Med* 2002;55:273-280.
- 37) Fettman MJ, Stanton CA, Banks LL, et al. Effects of neutering on body weight, metabolic rate and glucose tolerance of domestic cats. *Res Vet Sci* 1997;62:131-136.
- 38) Fidler IJ, Brodey RS. The biological behavior of canine mammary neoplasms. *J Amer Vet Med Assoc* 1967;151:1311-1318.
- 39) Flynn MF, Hardie EM, Armstrong PJ. Effect of ovariohysterectomy on maintenance energy requirement in cats. *J Amer Vet Med Assoc* 1996;209:1572-1581.
- 40) Fukuda S. Incidence of pyometra in colony-raised Beagle dogs. *Exp Anim* 2001;50:325-329.
- 41) Greenfield CL, Johnson AL, Schaeffer DJ. Frequency of use of various procedures, skills, and areas of knowledge among veterinarians in private small animal exclusive or predominant practice and proficiency expected of new veterinary school graduates. *J Amer Vet Med Assoc* 2004;224:1780-1787.
- 42) Greer KA, Canterbury SC, Murphy KE. Statistical analysis regarding the effects of height and weight on life span of the domestic dog. *Res Vet Sci* 2007;82:208-214.
- 43) Gregory SP, Parkinson TJ, Holt PE. Urethral conformation and position in relation to urinary incontinence in the bitch. *Vet Rec* 1992;131:167-170.
- 44) Gregory SP, Holt PE, Parkinson TJ, Wathes CM. Vaginal position and length in the bitch: Relationship to spaying and urinary incontinence. *J Sm Anim Prac* 1999;40:180-184.
- 45) Gunzel-Apel AR. Early castration of dogs and cats from the point of view of animal welfare. *Dtsch Tierarztl Woch* 1998;105:95-98.
- 46) Hahn KA, Vonderhaar MA, Teclaw RF. An epidemiological evaluation of 1202 dogs with testicular neoplasia. *J Vet Intern Med* 1992;6:121 [Abstract].
- 47) Hamilton JM, Else RW, Forshaw P. Oestrogen receptors in canine mammary tumours. *Vet Rec* 1977;101:258-260.

- 48) Hampe JF, Misdorp W. IX. Tumours and dysplasias of the mammary gland. *Bull World Hlth Org* 1974;50:111-133.
- 49) Hart BL. Effect of gonadectomy on subsequent development of age-related cognitive impairment in dogs. *J Amer Vet Med Assoc* 2001;219:51-56.
- 50) Hart BL, Barrett RE. Effects of castration on fighting, roaming, and urine spraying in adult male cats. *J Amer Vet Med Assoc* 1973;163:290-292.
- 51) Hart BL, Cooper L. Factors relating to urine spraying and fighting in prepubertally gonadectomized cats. *J Amer Vet Med Assoc* 1984;184:1255-1258.
- 52) Hart BL, Eckstein RA. The role of gonadal hormones in the occurrence of objectionable behaviours in dogs and cats. *Appl Anim Beh Sci* 1997;52:331-344.
- 53) Hayes HM, Milne KL, Mandel CP. Epidemiological features of feline mammary carcinoma. *Vet REc* 1981;108:476-479.
- 54) Head E, Moffat K, Das P, et al. Beta-amyloid deposition and tau phosphorylation in clinically characterized aged cats. *Neurobiol Aging* 2005;26:749-763.
- 55) Herron MA. The effect of prepubertal castration on the penile urethra of the cat. *J Amer Vet Med Assoc* 1972;160:208-211.
- 56) Herron MA. A potential consequence of prepuberal feline castration. *Fel Prac* 1971;1:17-19.
- 57) Holt PE. Urinary incontinence in the male and female dog or does sex matter? www.vin.com, 2004.
- 58) Holt PE, Thrusfield MV. Association in bitches between breed, size, neutering and docking, and acquired urinary incontinence due to incompetence of the urethral sphincter mechanism. *Vet Rec* 1993;133:177-180.
- 59) Hopkins SG, Schubert TA, Hart BL. Castration of adult male dogs: Effects on roaming, aggression, urine marking, and mounting. *J Amer Vet Med Assoc* 1976;168:1108-1110.
- 60) Houlton JEF, McGlennon NJ. Castration and physeal closure in the cat. *Vet Rec* 1992;131:466-467.
- 61) Haupt KA, Coren B, Hintz HF, Hilderbrant JE. Effect of sex and reproductive status on sucrose preference, food intake, and body weight of dogs. *J Amer Vet Med Assoc* 1979;174:1083-1085.
- 62) Howe LM. Surgical methods of contraception and sterilization. *Theriogenology* 2006;66:500-509.
- 63) Howe LM. Short-term results and complications of prepubertal gonadectomy in cats and dogs. *J Amer Vet Med Assoc* 1997;211:57-62.
- 64) Howe LM, Slater MR, Boothe HW, et al. Long-term outcome of gonadectomy performed at an early age or traditional age in cats. *J Amer Vet Med Assoc* 2000;217:1661-1665.
- 65) Howe LM, Slater MR, Boothe HW, et al. Long-term outcome of gonadectomy performed at an early age or traditional age in dogs. *J Amer Vet Med Assoc* 2001;218:217-221.
- 66) Howe LM. Prepubertal gonadectomy in dogs and cats – Part I. *Comp Cont Ed* 1999;21:103-111.
- 67) Howe LM. Prepubertal gonadectomy in dogs and cats – Part II. *Comp Cont Ed* 1999;21:197-201.
- 68) Janowsky JS. The role of androgens in cognition and brain aging in men. *Neurosci* 2006;138:1015-1020.

- 69) Johnson-Delaney CA. Ferret adrenal disease: 2006 perspective. *Exotic DVM* 2006;8:31-34.
- 70) Johnston SD, Root Kustritz MV, Olson PN. *Feline and canine theriogenology*, WB Saunders, Philadelphia, 2001:
- 71) Kaneene JB, Mostosky UV, Padgett GA. Retrospective cohort study of changes in hip joint phenotype of dogs in the United States. 1997;211:1542-1544.
- 72) Kass PH, New JC, Scarlett JM, Salman MD. Understanding animal companion surplus in the United States: Relinquishment of nonadoptables to animal shelters for euthanasia. *J Appl Anim Welf Sci* 2001;4:237-248.
- 73) Kealy RD, Olsson SE, Monti KL, et al. Effects of limited food consumption on the incidence of hip dysplasia in growing dogs. *J Amer Vet Med Assoc* 1992;201:857-863.
- 74) Keller GG, Corley EA. Canine hip dysplasia: Investigating the sex predilection and the frequency of unilateral CHD. *Vet Med* 1989;Dec: 1162-1166.
- 75) Kim HH, Yeon SC, Houpt KA, et al. Effects of ovariectomy on reactivity in German Shepherd dogs. *Vet J* 2006;172:154-159.
- 76) Knapp DW, Glickman NW, DeNicola DB, et al. Naturally-occurring canine transitional cell carcinoma of the urinary bladder. *Urologic Oncol* 2000;5:47-59.
- 77) Lawler DF, Evans RH, Reimers TJ, et al. Histopathologic features, environmental factors, and serum estrogen, progesterone, and prolactin values associated with ovarian phase and inflammatory uterine disease in cats. *Amer J Vet Res* 1991;52:1747-1753.
- 78) LeBoeuf BJ. Copulatory and aggressive behavior in the prepuberally castrated dog. *Horm Behav* 1970;1:127-136.
- 79) Leav I, Ling GV. Adenocarcinoma of the canine prostate gland. *Cancer* 1968;22:1329.
- 80) Lekcharoensuk C, Osborne CA, Lulich JP. Epidemiologic study of risk factors for lower urinary tract diseases in cats. *J Amer Vet Med Assoc* 2001;218:1429-1435.
- 81) Lipowitz AJ, Schwartz AS, Wilson GP. Testicular neoplasms and concomitant clinical changes in the dog. *J Amer Vet Med Assoc* 1973;163:1364-1368.
- 82) Lowseth LA, Gerlach RF, Gillett NA, Muggenburg BA. Age-related changes in the prostate and testes of the Beagle dog. *Vet Pathol* 1990;27:347-353.
- 83) Mackenzie AR, Hall T, Lo M-C, Whitmore WF. Influence of castration and sex hormones on size, histology and zinc content of canine prostate. *J Urol* 1963;89:864-874.
- 84) Manning AM, Rowan AN. Companion animal demographics and sterilization status: Results from a survey in four Massachusetts towns. *Anthrozoos* 1992;5:192-201.
- 85) Marmor M, Willeberg P, Glickman LT, et al. Epizootologic patterns of diabetes mellitus in dogs. *Amer J Vet Res* 1982;43:465-470.
- 86) Martin LJM, Siliart B, Dumon HJW, Nguyen P. Spontaneous hormonal variations in male cats following gonadectomy. *J Fel Med Surg* 2006;8:309-314.
- 87) Mason E. Obesity in pet dogs. *Vet Rec* 1970;86:612-616.
- 88) May C, Bennett D, Downham DY. Delayed physeal closure associated with castration in cats. *J Sm Anim Prac* 1991;32:326-328.

- 89) McCarthy MM, McDonald EH, Brooks PJ, Goldman D. An anxiolytic action of oxytocin is enhanced by estrogen in the mouse. *Phys Behav* 1997;60:1209-1215.
- 90) McNicholas WT, Wilkens BE, Blevins WE, et al. Spontaneous femoral capital physal fractures in adult cats: 26 cases (1996-2001). *J Amer Vet Med Assoc* 2002;221:1731-1736.
- 91) Michell AR. Longevity of British breeds of dog and its relationship with sex, size, cardiovascular variables and disease. *Vet Rec* 1996;145:625-629.
- 92) Misdorp W. Canine mammary tumours: Protective effect of late ovariectomy and stimulating effect of progestins. *Vet Quar* 1988;10:26-31.
- 93) Misdorp W, Hart AAM. Canine mammary cancer. II. Therapy and causes of death. *J Sm Anim Prac* 1979;20:395-404.
- 94) Mondelli F, Previde EP, Verga M, et al. The bond that never developed: Adoption and relinquishment of dogs in a rescue shelter. *J Appl Anim Welf Sci* 2004;7:253-266.
- 95) Moore GE, Burkman KD, Carter MN, Peterson MR. Causes of death or reasons for euthanasia in military working dogs: 927 cases (1993-1996). *J Amer Vet Med Assoc* 2001;219:209-214.
- 96) Moulton JE, Rosenblatt LS, Goldman M. Mammary tumors in a colony of Beagle dogs. *Vet Path* 1986;23:741-749.
- 97) Moulton JE, Taylor DON, Dorn CR, Andersen AC. Canine mammary tumors. *Path Vet* 1970;7:289-320.
- 98) Nassar R, Talboy J, Moulton C. Animal shelter reporting study 1990, Englewood CO, American Humane Assoc 1992:5.
- 99) National Council on Pet Population Study and Policy. National Shelter Census: 1994 Results. Fort Collins CO, 1994:1-2.
- 100) National Council on Pet Population Study and Policy. Shelter statistic survey, 1994-1997. www.petpopulation.org, 2007.
- 101) National Center for Health Statistics. Health, United States 2006: With chartbook on trends in the health of Americans, Hyattsville MD, 2006:176.
- 102) New JC, Salman MD, Scarlett JM, et al. Shelter relinquishment: Characteristics of shelter-relinquished animals and their owners compared with animals and their owners in US pet-owning households. *J Appl Anim Welf Sci* 2000;3:179-201.
- 103) New JC, Kelch WJ, Hutchison JM, et al. Birth and death rate estimates of cats and dogs in US households and related factors. *J Appl Anim Welf Sci* 2004;7:229-241.
- 104) Nguyen PG, et al. Effects of dietary fat and energy on body weight and composition after gonadectomy in cats. *Amer J Vet Res* 2004;65:1708-1713.
- 105) Nielsen JC, Eckstein RA, Hart BL. Effects of castration on problem behaviors in male dogs with reference to age and duration of behavior. *J Amer Vet Med Assoc* 1997;211:180-182.
- 106) Norris AM, Laing EJ, Valli VEO, et al. Canine bladder and urethral tumors: A retrospective study of 115 cases (1980-1985). *J Vet Intern Med* 1992;6:145-153.
- 107) Obradovich J, Walshaw R, Goulland E. The influence of castration on the development of prostatic carcinoma in the dog: 43 cases (1978-1985). *J Vet Intern Med* 1987;1:183-187.

- 108) O'Farrell V, Peachey E. Behavioural effects of ovariohysterectomy on bitches. *J Sm Anim Prac* 1990;31:595-598.
- 109) Olson PN. Early spay and neuter. Proceedings, North American Veterinary Conference, Orlando FL, 1997:25.
- 110) O'Shea JD. Studies on the canine prostate gland. I. Factors influencing its size and weight. *J Comp Path* 1962;72:321-331.
- 111) Osborne CA, Low DG, Perman V, et al. Neoplasms of the canine and feline urinary bladder: Incidence, etiologic factors, occurrence and pathologic features. *Amer J Vet Res* 1968;29:2041-2055.
- 112) Pabon JE, Li X, Lei ZM, et al. Novel presence of luteinising hormone / chorionic gonadotropin receptors in human adrenal glands. *J Clin Endocrin Metab* 1996;81:2397-2400.
- 113) Panciera DL. Hypothyroidism in dogs: 66 cases (1987-1992). *J Amer Vet Med Assoc* 1994;204:761-767.
- 114) Patronek GJ, Glickman LT, Beck AM, et al. Risk factors for relinquishment of dogs to an animal shelter. *J Amer Vet Med Assoc* 1996;209:572-581.
- 115) Patronek GJ, Glickman LT, Beck AM, et al. Risk factors for relinquishment of cats to an animal shelter. *J Amer Vet Med Assoc* 1996;209:582-588.
- 116) Perez Alenza MD, Tabanera E, Pena L. Inflammatory mammary carcinoma in dogs: 33 cases (1995-1999). *J Amer Vet Med Assoc* 2001;219:1110-1114.
- 117) Pollari FL, Bonnett BW, Bamsey SC, et al. Postoperative complications of elective surgeries in dogs and cats determined by examining electronic and paper medical records. *J Amer Vet Med Assoc* 1996;208:1882-1886.
- 118) Potter K, Hancock DH, Gallina AM. Clinical and pathologic features of endometrial hyperplasia, pyometra, and endometritis in cats: 79 cases (1980-1985). *J Amer Vet Med Assoc* 1991;198:1427-1431.
- 119) Priester WA, McKay FW. The occurrence of tumors in domestic animals. *Nat Canc Inst Monograph* 1980;54:169.
- 120) Prymak C, McKee LJ, Goldschmidt MH, Glickman LT. Epidemiologic, clinical, pathologic, and prognostic characteristics of splenic hemangiosarcoma and splenic hematoma in dogs: 217 cases (1985). *J Amer Vet Med Assoc* 1988;193:706-712.
- 121) Reichler IM, Welle M, Sattler U, et al. Comparative quantitative assessment of GnRH- and LH-receptor mRNA expression in the urinary tract of sexually intact and spayed female dogs. *Theriogenology* 2007;67:1134-1142.
- 122) Reichler IM, Pfeiffer E, Piche CA, et al. Changes in plasma gonadotropin concentrations and urethral closure pressure in the bitch during the 12 months following ovariectomy. *Theriogenology* 2004;62:1391-1402.
- 123) Reichler IM, Hung E, Jochle W, et al. FSH and LH plasma levels in bitches with differences in risk for urinary incontinence. *Theriogenology* 2005;63:2164-2180.
- 124) Reichler IM, Hubler M, Jochle W, et al. The effect of GnRH analogs on urinary incontinence after ablation of the ovaries in dogs. *Theriogenology* 2003;60:1207-1216.

- 125) Reisner IR. Dominance-related aggression of English springer spaniels: A review of 53 cases. *Appl Anim Behav Sci* 1993;37:83-84.
- 126) Robertson OH. Prolongation of the life span of Kokanee salmon (*Oncorhynchus nerka* kennerly) by castration before beginning of gonad development. *Proceedings, National Academies of Science USA* 1961;47:609-621.
- 127) Root MV, Johnston SD, Olson PN. The effect of prepuberal and postpuberal gonadectomy on radial physeal closure in male and female domestic cats. *Vet Rad US* 1997;38:42-47.
- 128) Root MV, Johnston SD, Johnston GR, Olson PN. The effect of prepuberal and postpuberal gonadectomy on penile extrusion and urethral diameter in the domestic cat. *Vet Rad US* 1996;37:363-366.
- 129) Root Kustritz MV. Elective gonadectomy in the cat. *Fel Prac* 1996;24:36-39.
- 130) Root Kustritz MV, Johnston SD, Lieberman LL. Availability of training for prepuberal gonadectomy at North American veterinary colleges. *J Amer Vet Med Assoc* 2000;216:1566-1567.
- 131) Root Kustritz MV. Determining the optimal age for gonadectomy of dogs and cats. *J Amer Vet Med Assoc* 2007; 231:1665 – 1675.
- 132) Rosenblatt JS, Aronson LR. The decline of sexual behavior in male cats after castration with special reference to the role of prior sexual experience. *Behavior* 1958;12:285-338.
- 133) Rosenblatt JS, Aronson LR. The influence of experience on the behavioural effects of androgen in prepuberally castrated male cats. *Anim Behav* 1958;6:171-182.
- 134) Rosenthal KL, Peterson ME, Quesenberry KE, et al. Hyperadrenocorticism associated with adrenocortical tumor or nodular hyperplasia of the adrenal gland in ferrets: 50 cases (1987-1991). *J Amer Vet Med Assoc* 1993;203:271-275.
- 135) Ru G, Terracini B, Glickman LT, et al. Related risk factors for canine osteosarcoma. *Vet J* 1998;156:31-39.
- 136) Sakata JT, Jupta A, Gonzalez-Lima F, Crews D. Heterosexual housing increases the retention of courtship behavior following castration and elevates metabolic capacity in limbic brain nuclei in male whiptail lizards, *Cnemidophorus inornatus*. *Horm Behav* 2002;42:263-273.
- 137) Salman MD, Hutchison J, Ruch-Gallie R, et al. Behavioral reasons for relinquishment of dogs and cats to 12 shelters. *J Appl Anim Welf Sci* 2000;3:93-106.
- 138) Salmeri KR, Bloomberg MS, Scruggs SL, Shille V. Gonadectomy in immature dogs: Effects on skeletal, physical and behavioral development. *J Amer Vet Med Assoc* 1991;198:1193-1203.
- 139) Salmeri KR, Olson PN, Bloomberg MS. Elective gonadectomy in dogs: A review. *J Amer Vet Med Assoc* 1991;198:1183-1192.
- 140) Scarlett JM, Salman MD, New JC, Kass PH. The role of veterinary practitioners in reducing dog and cat relinquishments and euthanasias. *J Amer Vet Med Assoc* 2002;220:306-311.
- 141) Scarlett JM, Salman MD, New JC, Kass PH. Reasons for relinquishment of companion animals in US animal shelters: Selected health and personal issues. *J Appl Anim Welf Sci* 1999;2:41-57.
- 142) Schneider R, Dorn CR, Taylor DON. Factors influencing canine mammary cancer development and postsurgical survival. *J Nat Canc Inst* 1969;43:1249-1261.

- 143) Seguin MA, Vaden SL, Altier C, et al. Persistent urinary tract infections and reinfections in 100 dogs (1989-1999). *J Vet Intern Med* 2003;17:622-631.
- 144) Shen V, Dempster DW, Birchman R, et al. Lack of changes in histomorphometric, bone mass, and biochemical parameters in ovari hysterectomized dogs. *Bone* 1992;13:311-316.
- 145) Shoemaker NJ, Schuurmans M, Moorman H, Lumeij JT. Correlation between age at neutering and age at onset of hyperadrenocorticism in ferrets. *J Amer Vet Med Assoc* 2000;216:195-197.
- 146) Slauterbeck JR, Pankratz K, Xu KT, et al. Canine ovari hysterectomy and orchiectomy increases the prevalence of ACL injury. *Clin Orthop* 2004;429:301-305.
- 147) Sloth C. Practical management of obesity in dogs and cats. *J Sm Anim Prac* 1992;33:178-182.
- 148) Sorenmo KU, Goldschmidt M, Shofer F, et al. Immunohistochemical characterization of canine prostatic carcinoma and correlation with castration status and castration time. *Vet Comp Oncol* 2003;1:48-56.
- 149) Spain CV, Scarlett JM, Houpt KA. Long-term risks and benefits of early-age gonadectomy in cats. *J Amer Vet Med Assoc* 2004;224:372-379.
- 150) Spain CV, Scarlett JM, Houpt KA. Long-term risks and benefits of early-age gonadectomy in dogs. *J Amer Vet Med Assoc* 2004;224:380-387.
- 151) Spain CV, Scarlett JM, Cully SM. When to neuter dogs and cats: A survey of New York state veterinarians' practices and beliefs. *J Amer Anim Hosp Assoc* 2002;38:482-488.
- 152) Stocklin-Gautschi NM, Hassig M, Reichler IM, et al. The relationship of urinary incontinence to early spaying in bitches. *J Rep Fert* 2001;Suppl 57:233-236.
- 153) Stubbs WP, Bloomberg MS, Scruggs SL, et al. Effects of prepubertal gonadectomy on physical and behavioral development in cats. *J Amer Vet Med Assoc* 1996;209:1864-1871.
- 154) Tarvin G, Patnaik A, Greene R. Primary urethral tumors in dogs. *J Amer Vet Med Assoc* 1978;172:931-933.
- 155) Teske E, Naan EC, VanDijk EM, et al. Canine prostate carcinoma: Epidemiological evidence of an increased risk in castrated dogs. *Molec Cell Endo* 2002;197:251-255.
- 156) Thrusfield MV, Holt PE, Muirhead RH. Acquired urinary incontinence in bitches: Its incidence and relationship to neutering practices. *J Sm Anim Prac* 1998;39:559-566.
- 157) Verstegen J, Onclin K. Etiopathogenesis, classification and prognosis of mammary tumors in the canine and feline species. *Proceedings, Society for Theriogenology, Columbus OH, 2003:230-238.*
- 158) Verstegen J, Onclin K. Mammary tumors in the queen. *Proceedings, Society for Theriogenology, Columbus OH, 2003:239-245.*
- 159) Ware WA, Hopper DL. Cardiac tumors in dogs: 1982-1995. *J Vet Intern Med* 1999;13:95-103.
- 160) Waters DJ, Shen S, Glickman LT. Life expectancy, antagonistic pleiotropy, and the testis of dogs and men. *The Prostate* 2000;43:272-277.
- 161) Weaver AD. Fifteen cases of prostatic carcinoma in the dog. *Vet Rec* 1981;109:71-75.
- 162) Whitehair JG, Vasseur PB, Willits NH. Epidemiology of cranial cruciate ligament rupture in dogs. *J Amer Vet Med Assoc* 1993;203:1016-1019.

163) Wright JC. Early-age gonadectomy and the behavior of kittens reared in a household setting. Proceedings, Midwest Veterinary Conference, Columbus OH, 2001:89-90.

164) Zirkin BR, Strandberg JD. Quantitative changes in the morphology of the aging canine prostate. Anat Rec 1984;208:207